



Role of Heparin during Pregnancy

Pedada Akhil *

Department of Medical Science, Kyrgyz-Russian Slavic University, Bishkek, Russia

Corresponding author email: akhilpedada410@gmail.com

Received on: 26-03-2021; Revised on: 09-04-2021; Accepted on: 16-04-2021

DESCRIPTION

Different kinds of heparin the sort we for the most part use is a Low Molecular Weight Heparin (LMWH) like Enoxaparin (otherwise called Clexane) or Dalteparin.

Venous apoplexy is a condition wherein blood coagulation (clots) structures in a vein (venous apoplexy). Most regularly, venous apoplexy happens in the "profound veins" in the legs, thighs or pelvis and this is known as a profound vein apoplexy or a DVT. A DVT can restrict blood course through the vein, causing expanding and torment, yet there is a danger that it can sever and go through the circulation system. This is called an embolus. On the off chance that the embolus lodges in the lung this is known as a pneumonic embolus (PE). A PE may mess breathing up, chest torment and hacking up blood. During pregnancy and the Heparin is an anticoagulant drug. This implies that is hinders blood coagulating. There are prompt period after birth, the danger of venous apoplexy is expanded.

Role of heparin

All in all, LMWHs are utilized widely during pregnancy and puerperium in Greece for VTE treatment and prophylaxis and for an assortment of different signs also. This examination is the main public overview in regards to LMWHs use in pregnancy and exhibits the wellbeing and viability of the medication in a high danger pregnancy setting. It is of incredible worry that over remedy of LMWHs in pregnant ladies with a background marked by GVCs, in spite of the absence of strong logical information, addresses the current clinical practice all through our country and past. The examination restrictions are its non-controlled, review nature, and

the heterogeneity of signs for LMWHs organization. The unseemly utilization of these medications ought to be forestalled by setting up and executing symptomatic and restorative rules and giving the essential schooling to medical care experts. To unmistakably characterize the job of LMWHs in pregnancy, past thromboprophylaxis, enormous imminent examinations are required, which could be founded on the finishes of this investigation.

The three most basic reasons for ladies to begin heparin during pregnancy or after the birth are:

- An expanded danger of a coagulation framing during pregnancy or after birth, for example after some cesarean births or in the event that you are stable for a while.
- A coagulation emerging during this or a past pregnancy.
- Effectively on long haul anticoagulation, for example repetitive apoplexy.

CONCLUSION

All in all, LMWHs are utilized widely during pregnancy and puerperium in Greece for VTE treatment and prophylaxis and for an assortment of different signs also. This examination is the main public overview in regards to LMWHs use in pregnancy and exhibits the wellbeing and viability of the medication in a high danger pregnancy setting. It is of incredible worry that over remedy of LMWHs in pregnant ladies with a background marked by GVCs, in spite of the absence of strong logical information, addresses the current clinical practice all through our country and past. The examination restrictions are its non-controlled, review nature, and

the heterogeneity of signs for LMWHs organization. The unseemly utilization of these medications ought to be forestalled by setting up and executing symptomatic and restorative rules and giving the essential schooling to medical care experts. To unmistakably characterize the job of LMWHs in pregnancy, past thromboprophylaxis, enormous imminent examinations are required, which could be founded on the finishes of this investigation.

REFERENCE

- [1] Warkentin TE, Kelton JG. A 14-year study of heparin-induced thrombocytopenia. *Am J Med* .1996; 101:502-507.
- [2] Warkentin TE, Levine MN, Hirsh J, et al. Heparin-induced thrombocytopenia in patients treated with low-molecular-weight heparin or unfractionated heparin. *N Engl J Med* 1995; 332:1330-1335.
- [3] Kaplan FS. Prevention and management of osteoporosis. *Clin Symp* 1995; 47:2-32.
- [4] Shefras J, Farquharson RG. Bone density studies in pregnant women receiving heparin. *Eur J Obstet Gynecol Reprod Biol* 1996; 65:171-174.
- [5] Meytes D, Ayalon H, Virag I, Weisbort Y, Zakut H. Heparin-induced thrombocytopenia and recurrent thrombosis in pregnancy: a case report. *J Reprod Med* 1986; 31:993-996.